

SUMMARY OF AUSTRALIAN INDIGENOUS HEALTH, FEBRUARY 2005

Australian Indigenous Health *InfoNet*

Introduction

This summary includes facts about common health problems and risk factors among Australian Indigenous people. More detailed information about the health of Indigenous people, associated social and economic circumstances, and health risk factors, is available from the *HealthInfoNet's* website (www.healthinfonet.ecu.edu.au).

Indigenous Population

There were around 483,990 Indigenous people living in Australia in 2004 (around 432,560 Aboriginal people, 30,740 Torres Strait Islanders, and 20,690 people of both Aboriginal and Torres Strait Islander descent) [1, 2]. Indigenous people comprise around 2.4% of the total Australian population. Most Indigenous people live in New South Wales (NSW), followed by Queensland (Qld), Western Australia (WA), and the Northern Territory (NT). The NT has the highest percentage of Indigenous people among its population and Victoria (Vic) the lowest. Most Torres Strait Islander people live in Queensland, with NSW the only other State with a large number of Torres Strait Islanders.

The Indigenous population is much younger overall than the non-Indigenous population [1]. According to the 2001 Australian Census, about 40 out of 100 Indigenous people were aged less than 15 years, compared with 20 out of 100 non-Indigenous people. About 3 out of 100 Indigenous people were aged 65 years or over, compared with 10 out of 100 non-Indigenous people.

Births

In 2003, there were 11,740 births registered in Australia where one or both parents were Indigenous (five out of every one hundred births) [3]. Overall, Indigenous women had more children and had them at younger ages than did non-Indigenous women.

Based on the pattern of births in recent years, Indigenous women would have, on average, around 2.15 births in their lifetime, compared with less than 1.8 births for non-Indigenous women [3]. More than 52 out of 100 Indigenous women are 24 years or younger when they have their babies, compared with less than 18 out of 100 non-Indigenous women. More than 21 in 100 Indigenous mothers are teenagers, compared with less than four in 100 non-Indigenous mothers.

On average, babies born to Indigenous women in recent years have weighed around 200 grams less than those born to non-Indigenous women [4]. Babies born to Indigenous women are more than twice as likely to be of low birth weight (less than

2,500 grams) than are those born to non-Indigenous women [4]. (Low birth weight can increase the risk of health problems.)

Deaths

Indigenous people are much more likely to die before they are old than people in the rest of the Australian population [5]. Estimates from the Australian Bureau of Statistics (ABS) indicate that at birth an Indigenous male born in the period 1996–2001 could be expected to live to 59 years, which is around 17 years less than a male in the total population at that time (who had a life expectancy of 76.5 years). In the same period, an Indigenous female could be expected to live to 65 years, which is around 17 years less than a woman in the total population (82 years).

In 2003, there were 2,079 people who died and were identified as Indigenous [5]. Many Indigenous deaths are incorrectly identified as non-Indigenous—the actual number of Indigenous deaths is likely to be around 3,600.

Death rates relate the numbers of deaths to the total numbers of people. After taking account of the facts that the Indigenous population is much younger overall than the non-Indigenous population and that many Indigenous deaths are not identified as such, the death rates for Indigenous males and females are likely to be around four times higher than those of their non-Indigenous counterparts [6].

Indigenous babies are more likely to die in their first year than non-Indigenous babies [5]. In 2001–03, the infant mortality rate for Indigenous babies was highest in WA (16 babies died out of 1,000 births) and the NT (16 babies died out of 1,000 births) and lowest in NSW (9 babies died out of 1,000 births). (The rate for the total Australian population is around 5 deaths per 1,000 births.)

In 2000–02 the leading causes of death for Indigenous people living in Qld, WA, SA and the NT were: cardiovascular disease (including heart disease and strokes); injuries (including transport accidents, self-harm and assault); cancer; respiratory diseases; and diabetes [7]. (More information about these causes of death is provided below).

Specific Health Conditions

Cardiovascular Disease

More than one in four deaths registered as Indigenous in recent years were caused by cardiovascular disease [7, 8]. Indigenous people are much more likely to die from cardiovascular disease than other Australians at any age, and particularly in younger age groups. The cardiovascular disease death rate among Indigenous people aged between 25 and 54 years is at least eight to ten times, and possibly as high as 15 times, that of other Australians [9].

In the 2001 National Health Survey (NHS), about one in ten Indigenous people reported having a long-term cardiovascular condition, particularly in older age groups [10]. The most commonly reported condition was hypertension (high blood pressure).

Cancer

It is not known just how many Indigenous people develop cancer, but notification rates for new cases have been lower for Indigenous people than for non-Indigenous people in recent years [9]. On the other hand, death rates for people living in Qld, WA and the NT are generally higher for Indigenous people than for non-Indigenous people (these rates take account of the fact that the Indigenous population is much younger overall than the non-Indigenous population) [9, 11].

The leading causes of Indigenous cancer deaths include cancers of the digestive organs and lung cancer [8, 12]. Indigenous people have higher rates of smoking-related cancers than non-Indigenous people [9, 11, 12]. Indigenous women have higher rates of cervical cancer than non-Indigenous women, but lower rates of breast cancer.

The fact that Indigenous people are more likely than non-Indigenous people to die from cancer could be because the cancers they develop (such as cancers of the lung and liver) are more likely to be fatal, or that the stage of cancer may be more advanced by the time it is recognised [9, 11].

Diabetes

Diabetes is a major health problem among Indigenous people, but it is difficult to know just how many Indigenous people have the disease. The best evidence suggests that diabetes is between two and four times more common among Indigenous people than among non-Indigenous people [13]. Indigenous people are likely to be diagnosed with diabetes at a much lower age than non-Indigenous people [14]. Deaths from diabetes are much more common for Indigenous people than for non-Indigenous people [8]. In recent years in Qld, WA and the NT, diabetes accounted for 11 times as many deaths as expected for Indigenous males and 18 times as many deaths as expected for Indigenous females (based on total Australian male and female rates, but, as with other estimated death rates, the actual difference is likely to be up to 30% greater) [9].

In the 2001 National Health Survey, five out of every one hundred Indigenous people reported that they had diabetes as a 'long-term health condition' [10]. Indigenous people living in remote areas were more likely to report having diabetes than Indigenous people in other areas. (It should be noted that for every person who reports in surveys that they have diabetes it is likely that there is another person who doesn't know they have the disease [15].)

Renal Disease

Renal disease, which affects the kidneys, has only recently been fully recognised as a serious public health threat to Indigenous people. End-stage renal disease (ESRD) occurs when the kidneys are no longer able to function. Rates of ESRD are much higher for Indigenous people than they are for non-Indigenous people across most of the country, and particularly in remote areas where they are up to 30 times higher [16]. Death rates from chronic kidney disease for people living in Qld, WA, SA and

the NT in 1999–2001 were eight times higher for Indigenous people than for non-Indigenous people [9].

Dialysis (the usual treatment for ESRD—where the work of the kidneys is done artificially) accounted for more than one-third of all hospital admissions among Indigenous people in 2002–03 (many of these involved repeat admissions for the same people, some on an almost daily basis) [17]. Indigenous people were seven-and-a-half times more likely to be hospitalised for dialysis than non-Indigenous people. In recent years, almost half of all Indigenous ESRD patients have come from regions without dialysis or transplant facilities, and around one in six from regions with only satellite dialysis facilities.

Injury

Indigenous people are more likely to die from transport accidents, intentional self-harm and assault than other Australians [8]. Injury was the second most common cause of death for Indigenous males living in Qld, WA, SA and the NT in 2000–02, and the fourth most common cause of death for Indigenous females—rates were around three times those of the total Australian populations (the actual difference is likely to be up to 30% greater) [7].

Across Australia in 2002–03, Indigenous people were around twice as likely as other Australians to be admitted to hospital for injuries [17]. Assault was the leading cause of hospitalisation as a result of injury for both Indigenous males and females, followed by accidental falls. Other common causes of hospitalisation for injury were transport accidents (particularly for Indigenous males), complications of medical and surgical care, and intentional self-harm. Hospitalisation rates for injury among Indigenous people were higher than those among non-Indigenous people in nearly every age group.

Respiratory Disease

About one in eleven of all deaths registered as Indigenous in recent years were due to a respiratory disorder [7, 8]. For Indigenous people living in Qld, WA, SA and the NT in 2000–02, deaths from these disorders were around four times more common than for non-Indigenous people (the actual difference is likely to be up to 30% greater) [7]. In the 35–44 years age group, the death rates for Indigenous people were around 20 times higher for males and 10 times higher for females than for other Australian males and females.

In recent years, Indigenous people were more likely to be hospitalised for respiratory disease than non-Indigenous people, particularly in infancy and early childhood [17].

Communicable Diseases

It is compulsory for some communicable diseases to be notified. The States and Territories collect the information, which is then collated and published by the National Notifiable Disease Surveillance System. Indigenous status is often not reported in notifications, and only the information from WA, SA and the NT is believed to be complete enough for publication by Indigenous status [9]. Information

for Australian HIV/AIDS cases relates to all States and Territories and is collated and published by the National Centre in HIV Epidemiology and Clinical Research [18].

Recent information about communicable diseases includes:

- tuberculosis – the rate of newly diagnosed cases for Indigenous people in 2003 was 10 times the rate for Australian-born non-Indigenous people [19];
- *Haemophilus influenzae* type B—the notification rate in 2000–02 for Indigenous children aged less than 5 years was 14 times that for the total Australian population in that age group [20];
- meningococcal infection—the notification rate for Indigenous people living in NSW, WA, SA and the NT in 2000–02 was around twice that of the total population of those States [20];
- syphilis and gonorrhoea—notification rates for Indigenous people living in WA, SA and the NT in 2003 were between 50 and 100 times higher than those for the total Australian population [18]; and
- HIV/AIDS—notification rates for HIV infection and AIDS are similar for the Indigenous population and the non-Indigenous population [18].

Eye Health

Eye conditions that affect the Indigenous population include refractive error (requiring glasses for correction), cataract (clouding of the lens), trachoma (a bacterial infection that can lead to blindness if untreated) and diabetic retinopathy (damage to the retina, at the back of the eye, caused by diabetes). There has been progress in the eye health of Indigenous people, but many are still more likely than non-Indigenous people to suffer from preventable conditions [21]. The frequency and severity of trachoma, for example, have decreased generally, but the infection is still quite common among Indigenous children living in some remote parts of the country.

The eye health of many Indigenous people is limited also by their difficulty in accessing specialised ophthalmological or optometrist services (because they are not available where they live, or are not culturally appropriate, or they are too expensive).

Ear Conditions

According to the 2001 National Health Survey, almost one in five Indigenous people have some degree of hearing loss, compared with around one in seven non-Indigenous people [10]. This higher frequency of hearing loss, reported for all age groups except people aged 55 years or older, reflects mainly the much higher levels of otitis media (middle ear disease) that Indigenous people experience in their childhood years. The actual levels vary considerably—geographically and over time—but it was estimated that in the mid 1990s around one in seven Indigenous children under 10 years of age had a perforated eardrum [22]. This level puts Indigenous children within the World Health Organization's 'extremely high risk' category. Disease of the middle ear can cause permanent hearing loss that limits life opportunities, particularly in education and in employment [23, 24].

Oral Health

The oral health of Indigenous people generally is not as good as that of other Australians. In contrast to the situation for young non-Indigenous children, whose oral health has improved in recent years, that of young Indigenous children has generally declined [25]. (There may be differences across the country, however, as Indigenous children in remote areas appear to have better oral health than those in urban areas.) Partly because the overall level of dental care is lower for Indigenous people than for non-Indigenous people, their oral health deteriorates with age, contributing to higher frequencies of periodontal (gum) disease and tooth loss.

Skin Infections and Infestations

Skin infections, which are more common for Indigenous people than for non-Indigenous people, are often the result of poor living conditions. The most common skin infections affecting Indigenous people are scabies (caused by a mite) and streptococcal pyoderma (a bacterial infection) [26]. Scabies, in particular, is a problem in many remote Indigenous communities where up to half the children may be infected.

Skin diseases cause very few deaths directly, but they can be linked with serious complications. They did, however, account for around 1 out of every 40 hospital separations in 2002–03 for patients identified as Indigenous, at a rate around two-and-a-half that of non-Indigenous people [17].

Factors Contributing to Ill-health

Indigenous people generally experience more risk factors for ill-health than do other Australians. Contributing generally to the poor health status of many Indigenous people are: social factors such as dispossession, dislocation and discrimination; disadvantages in education, housing, income and employment; and physical environmental factors. These social, economic and environmental disadvantages underlie specific health risk factors (such as smoking, obesity, physical inactivity and high blood pressure), and often contribute to lack of access to good quality health care.

Given the importance of these factors, substantial improvements in Indigenous health status are unlikely to be achieved without improvements in the overall circumstances of Indigenous people.

Summary

Indigenous people remain the least healthy sub-population in Australia, and there is evidence that the difference between Indigenous and non-Indigenous health, at least measured by mortality, has widened in recent years [27].

The reasons why the health status of Indigenous people remains much worse than that of non-Indigenous people are complex, but represent a combination of general factors—such as education, employment, income and socioeconomic status—and

EXPOSURE DRAFT FOR CONSULTATION

factors more specific to the health sector. As the Australian health ministers noted in their introduction to the 2003 *National Strategic Framework for Aboriginal and Torres Strait Islander Health*, achievement of substantial improvements in Indigenous health will depend on long-term collaborative approaches involving Indigenous leaders and communities, the health and non-health sectors, and all levels of government [28].

Within the health sector, there is a need for further improvement in: health advancement programs; identification of health conditions before they become serious; and expansion of primary health care services. To achieve long-term health benefits, funding needs to be directed to a wide range of preventive and clinical services. Funding should take account of the fact that mainstream services may not be accessible for many Indigenous people who may also have difficulty in accessing Medicare and pharmaceutical benefits.

However, without substantial reductions in the overall disadvantages experienced by many Indigenous people, even fully committed approaches within the health sector will have a limited impact on achieving major improvements in Indigenous health status.

References

- 1 Australian Bureau of Statistics (2003) *Australian Demographic Statistics Quarterly: 2001. March quarter 2003*. (Cat no. 3101.0) Canberra: Australian Bureau of Statistics
- 2 Australian Bureau of Statistics (2004) *Experimental estimates and projections, Aboriginal and Torres Strait Islander Australians*. (Cat No. 3238.0) Canberra: Australian Bureau of Statistics
- 3 Australian Bureau of Statistics (2004) *Births, Australia 2003*. (3301.0) Canberra: Australian Bureau of Statistics
- 4 Laws PJ, Sullivan EA (2004) *Australia's mothers and babies 2001*. (AIHW cat. no. PER 25) Sydney: AIHW National Perinatal Statistics Unit
- 5 Australian Bureau of Statistics (2004) *Deaths Australia 2003*. (Cat no. 3302.0) Canberra: Australian Bureau of Statistics
- 6 Thomson N, Ali M (2003) Births, deaths, and hospitalisation. In: Thomson N, ed. *The health of Indigenous Australians*. South Melbourne: Oxford University Press:44-74
- 7 Australian Institute of Health and Welfare (2004) *Australia's health 2004: the ninth biennial report of the Australian Institute of Health and Welfare*. Canberra: Australian Institute of Health and Welfare
- 8 Australian Bureau of Statistics (2003) *Causes of Death 2002*. (Cat no. 3303.0) Canberra: Australian Bureau of Statistics
- 9 Australian Bureau of Statistics, Australian Institute of Health and Welfare (2003) *The health and welfare of Australia's Aboriginal and Torres Strait Islander peoples 2003*. (ABS Cat no.4704.0, AIHW Cat no. IHW11) Canberra: Australian Bureau of Statistics
- 10 Australian Bureau of Statistics (2002) *National Health Survey: Aboriginal and Torres Strait Islander results, Australia 2001*. (ABS Catalogue no. 4715.0) Canberra: Australian Bureau of Statistics
- 11 Kirov E, Francis J, Thomson N (2003) Cancer. In: Thomson N, ed. *The health of Indigenous Australians*. South Melbourne: Oxford University Press:207-223
- 12 Condon JR, Barnes T, Cunningham J, Armstrong BK (2004) Long-term trends in cancer mortality for Indigenous Australians in the Northern Territory. *Medical Journal of Australia*;180(10):504-511
- 13 de Courten M, Hodge A, Dowse G, King I, Vickery J, Zimmet P (1998) *Review of the epidemiology, aetiology, pathogenesis and preventability of diabetes in Aboriginal and Torres Strait Islander populations*. Canberra: Commonwealth Department of Health and Family Services

EXPOSURE DRAFT FOR CONSULTATION

- 14 Couzos S, Metcalf S, Murray R, O'Rourke S (1998) *Systematic review of existing evidence and primary care guidelines on the management of non- insulin dependent diabetes in Aboriginal and Torres Strait Islander populations*. Canberra: Office for Aboriginal and Torres Strait Islander Health Services, Commonwealth Department of Health and Family Services
- 15 Dunstan D, Zimmet P, Welborn T, et al. (2001) *Diabetes and associated disorders in Australia, 2000: the accelerating epidemic*. Melbourne: International Diabetes Institute
- 16 Cass A, Cunningham J, Wang Z, Hoy W (2001) Regional variation in the incidence of end-stage renal disease in Indigenous Australians. *Medical Journal of Australia*;175:24-27
- 17 Australian Institute of Health and Welfare (2004) *Australian hospital statistics 2002-03*. (AIHW cat. no. HSE 32) Canberra: Australian Institute of Health and Welfare
- 18 National Centre in HIV Epidemiology and Clinical Research (2004) *2004 annual surveillance report: HIV/AIDS, viral hepatitis and sexually transmissible infections in Australia*. Sydney: National Centre in HIV Epidemiology and Clinical Research
- 19 Li J, Roche P, Spencer J, and the National Tuberculosis Advisory Committee (2004) Tuberculosis notifications in Australia, 2003. *Communicable Diseases Intelligence*;28(4):464-473
- 20 Menzies R, McIntyre P, Beard F (2004) Vaccine preventable diseases and vaccination coverage in Aboriginal and Torres Strait Islander people, Australia, 1999 to 2002. *Communicable Diseases Intelligence*;28(2):127-159
- 21 Taylor HR (1997) *Eye health in Aboriginal and Torres Strait Islander communities: the report of a review commissioned by the Commonwealth Minister for Health and Family Services, the Hon. Michael Wooldridge, MP*. Canberra: Commonwealth Department of Health and Family Services
- 22 Morris P (1998) A systematic review of clinical research addressing the prevalence, aetiology, diagnosis, prognosis and therapy of otitis media in Australian Aboriginal children [review]. *Journal of Paediatrics and Child Health*;34(6):487-497
- 23 Couzos S, Metcalf S, Murray R (2003) Ear health. In: Couzos S, Murray R, eds. *Aboriginal primary health care: an evidence-based approach*. 2nd ed. South Melbourne: Oxford University Press:193-250
- 24 Burrow S, Thomson N (2003) Ear disease and hearing loss. In: Thomson N, ed. *The health of Indigenous Australians*. South Melbourne: Oxford University Press:247-272
- 25 Harford J, Spencer J, Roberts-Thomson K (2003) Oral health. In: Thomson N, ed. *The health of Indigenous Australians*. South Melbourne: Oxford University Press:313-338
- 26 Currie BJ, Carapetis JR (2000) Skin infections and infestations in Aboriginal communities in northern Australia. *Australasian Journal of Dermatology*;41(3):139-145
- 27 Ring IT, Firman D (1998) Reducing Indigenous mortality in Australia: lessons from overseas. *Medical Journal of Australia*;169:528-533
- 28 National Aboriginal and Torres Strait Islander Health Council (2003) *National Strategic Framework for Aboriginal and Torres Strait Islander Health: framework for action by governments*. Canberra: National Aboriginal and Torres Strait Islander Health Council

AUSTRALIAN INDIGENOUS HEALTHINFONET

The Australian Indigenous Health*InfoNet*'s mission is to contribute to improvements in Indigenous health by making relevant, high quality knowledge and information easily accessible to policy makers, health service providers, program managers, clinicians, researchers and the general community.

The Health*InfoNet* addresses this mission by undertaking research into various aspects of Indigenous health and disseminates the results (and other relevant knowledge and information) mainly via its Internet site (www.healthinfonet.ecu.edu.au). The Health*InfoNet*'s research mainly involves analysis and synthesis of data and other information obtained from academic, professional, government and other sources, but it also undertakes some primary data collection and analysis.

The Health*InfoNet* is a world leader in knowledge transfer, the area of research which aims at transferring the results of pure and applied research into practice. In this research, the Health*InfoNet* addresses the knowledge needs of a wide range of potential users. These include policy makers, health service providers, program managers, clinicians and other health professionals (including Indigenous health workers), and researchers. The Health*InfoNet* also provides easy-to-read and summarised material for students and the general community.